




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 471/14, A61K 31/44	A1	(11) International Publication Number: WO 00/26217 (43) International Publication Date: 11 May 2000 (11.05.00)
(21) International Application Number: PCT/EP99/08227 (22) International Filing Date: 29 October 1999 (29.10.99) (30) Priority Data: 98120834.1 3 November 1998 (03.11.98) EP (71) Applicant (for all designated States except US): BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH [DE/DE]; Byk-Gulden-Str. 2, D-78467 Konstanz (DE). (72) Inventors (for all designated States except CA US): GRUNDLER, Gerhard; Meersburger Str. 4, D-78464 Konstanz (DE). POSTIUS, Stefan; Austr. 4b, D-78467 Konstanz (DE). SIMON, Wolfgang-Alexander; Schubert-str. 17, D-78464 Konstanz (DE). KROMER, Wolfgang; Alpenstrasse 2, D-78464 Konstanz (DE). (72) Inventor; and (75) Inventor/Applicant (for US only): SENN-BILFINGER, Jörg [DE/DE]; Säntisstrasse 7, D-78464 Konstanz (DE). (74) Common Representative: BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH; Byk-Gulden-Str. 2, D-78467 Konstanz (DE).		(81) Designated States: AE, AL, AU, BA, BG, BR, CA, CN, CZ, EE, GE, HR, HU, ID, IL, IN, JP, KR, LT, LV, MK, MX, NO, NZ, PL, RO, SG, SI, SK, TR, UA, US, VN, YU, ZA, ZW, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>

(54) Title: IMIDAZONAPHTHYRIDINES**(57) Abstract**

Compounds of formula (I), in which the substituents are as defined in the description are suitable for preventing and treating gastrointestinal disorders.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Imidazonaphthyridines**Field of application of the invention**


The invention relates to novel compounds which are used in the pharmaceuticals industry as active compounds for preparing medicaments.

Known technical background

US Patent 4,468,400 describes tricyclic imidazo[1,2-a]pyridines having different ring systems fused to the imidazopyridine skeleton, which compounds are said to be suitable for treating peptic ulcer disorders. The International Patent Application WO98/42707 discloses tetrahydroimidazonaphthyridines having a very particular substitution pattern, which compounds are likewise said to be suitable for treating gastrointestinal disorders.

Description of the invention

The invention provides compounds of the formula I



in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl or hydroxy-1-4C-alkyl,

R3 is hydrogen or halogen,

one of the substituents R4a and R4b is hydrogen and the other is hydrogen, hydroxyl, 1-4C-alkoxy, 1-4C-alkoxy-1-4C-alkoxy, 1-4C-alkylcarbonyloxy or the radical R4', or R4a and R4b together are O (oxygen),

where

R4' is -O-(CH₂)_m-S(O)_n-R9,

$-\text{S(O)}_n--(\text{CH}_2)_m-\text{OH},$
 $-\text{S(O)}_n--(\text{CH}_2)_m-\text{O-R}_9,$
 $-\text{S(O)}_n--(\text{CH}_2)_m-\text{S(O)}_p-\text{R}_9,$
 $-\text{O-alk1-S(O)}_n-\text{R}_9,$
 $-\text{S(O)}_n-\text{R}_9,$
 $-\text{S(O)}_n-\text{alk1-OH},$
 $-\text{S(O)}_n-\text{alk1-O-R}_9$ or
 $-\text{S(O)}_n-\text{alk1-S(O)}_p-\text{R}_9,$

in which

R₉ is 1-7C-alkyl, halo-1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl or di-1-4C-alkylamino-1-4C-alkyl, Ar or Ar-1-4C-alkyl, where Ar is phenyl or substituted phenyl having one, two or three identical or different substituents selected from the group consisting of 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkylcarbonyl, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl, difluoromethoxy, trifluoromethoxy, amino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino and nitro,

alk1 is 1-4C-alkyl-, hydroxyl-, oxo-, carboxyl-, halogen-, amino-, 1-4C-alkoxycarbonylamino- or phenyl-substituted 2-7C-alkylene or 3-4C-alkenylene,

m is an integer from 2 to 7,

n is the number 0, 1 or 2 and

p is the number 0, 1 or 2,

one of the substituents R_{5a} and R_{5b} is hydrogen and the other is hydrogen, hydroxyl, 1-4C-alkoxy, 1-4C-alkoxy-1-4C-alkoxy, 1-4C-alkylcarbonyloxy or the radical R_{5'}, or R_{5a} and R_{5b} together are O (oxygen),

where

R_{5'} is $-\text{O}-(\text{CH}_2)_q-\text{S(O)}_r-\text{R}_{10},$
 $-\text{S(O)}_r-(\text{CH}_2)_q-\text{OH},$
 $-\text{S(O)}_r-(\text{CH}_2)_q-\text{O-R}_{10},$
 $-\text{S(O)}_r-(\text{CH}_2)_q-\text{S(O)}_r-\text{R}_{10},$
 $-\text{O-alk2-S(O)}_r-\text{R}_{10},$
 $-\text{S(O)}_r-\text{R}_{10},$
 $-\text{S(O)}_r-\text{alk2-OH},$
 $-\text{S(O)}_r-\text{alk2-O-R}_{10}$ or
 $-\text{S(O)}_r-\text{alk2-S(O)}_r-\text{R}_{10},$

in which

R₁₀ is 1-7C-alkyl, halo-1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl or di-1-4C-alkylamino-1-4C-alkyl, Ar or Ar-1-4C-alkyl, where Ar is phenyl or substituted phenyl having one, two or three identical or different substituents selected from the group consisting of 1-4C-alkyl, 1-4C-alkoxy,

1-4C-alkylcarbonyl, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl, difluoromethoxy, trifluoromethoxy, amino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino and nitro,

alk2 is 1-4C-alkyl-, hydroxyl-, oxo-, carboxyl-, halogen-, amino-, 1-4C-alkoxycarbonylamino- or phenyl-substituted 2-7C-alkylene or 3-4C-alkenylene,

q is an integer from 2 to 7,

r is the number 0, 1 or 2 and

t is the number 0, 1 or 2,

R6 is hydrogen, halogen, 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or trifluoromethyl,

R7 is hydrogen, halogen, 1-4C-alkyl or 1-4C-alkoxy and

R8 is hydrogen or 1-4C-alkyl,

and their salts,

where one of the substituents R4a and R4b has to be R4' and/or one of the substituents R5a and R5b has to be R5'.

1-4C-Alkyl represents straight-chain or branched alkyl radicals having 1 to 4 carbon atoms. Examples which may be mentioned are the butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl, ethyl and methyl radicals.

Hydroxy-1-4C-alkyl represents the abovementioned 1-4C-alkyl radicals, substituted by a hydroxyl group. Examples which may be mentioned are the hydroxymethyl, the 2-hydroxyethyl and the 3-hydroxypropyl radicals.

For the purpose of the invention, halogen is bromine, chlorine and fluorine.

1-4C-Alkoxy represents radicals which, in addition to the oxygen atom, contain a straight-chain or branched alkyl radical having 1 to 4 carbon atoms. Examples which may be mentioned are the butoxy, isobutoxy, sec-butoxy, tert-butoxy, propoxy, isopropoxy and preferably the ethoxy and methoxy radicals.

1-4C-Alkoxy-1-4C-alkoxy represents one of the abovementioned 1-4C-alkoxy radicals which is substituted by a further 1-4C-alkoxy radical. Examples which may be mentioned are the radicals 2-(methoxy)ethoxy($\text{CH}_3\text{-O-CH}_2\text{-CH}_2\text{-O-}$) and 2-(ethoxy)-ethoxy($\text{CH}_3\text{-CH}_2\text{-O-CH}_2\text{-CH}_2\text{-O-}$).

1-4C-Alkylcarbonyl represents a radical which, in addition to the carbonyl group, contains one of the abovementioned 1-4C-alkyl radicals. An example which may be mentioned is the acetyl radical.

1-4C-Alkylcarbonyloxy represents a carbonyloxy group to which is attached one of the abovementioned 1-4C-alkyl radicals. An example which may be mentioned is the acetoxy radical ($\text{CH}_3\text{CO-O-}$).

1-7C-Alkyl represents straight-chain or branched alkyl radicals having 1 to 7 carbon atoms. Examples which may be mentioned are the heptyl, isoheptyl (5-methylhexyl), hexyl, isoheptyl(4-methylpentyl), neoheptyl(3,3-dimethylbutyl), pentyl, isopentyl(3-methylbutyl), neopentyl(2,2-dimethylpropyl), butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl, ethyl and methyl radicals.

Halo-1-4C-alkyl represents one of the abovementioned 1-4C-alkyl radicals which is substituted by one of the abovementioned halogen atoms. An example which may be mentioned is the 3-chloropropyl radical.

1-4C-Alkoxy-1-4C-alkyl represents one of the above-mentioned 1-4C-alkyl radicals which is substituted by one of the abovementioned 1-4C-alkoxy radicals. Examples which may be mentioned are the methoxymethyl, the methoxyethyl and the butoxyethyl radicals.

Carboxy-1-4C-alkyl represents, for example, the carboxy-methyl ($-\text{CH}_2\text{COOH}$) or the carboxyethyl ($-\text{CH}_2\text{CH}_2\text{COOH}$) radicals.

1-4C-Alkoxy carbonyl represents a carbonyl group to which is attached one of the abovementioned 1-4C-alkoxy radicals. Examples which may be mentioned are the methoxycarbonyl ($\text{CH}_3\text{O-C(O)-}$) and the ethoxycarbonyl ($\text{CH}_3\text{CH}_2\text{O-C(O)-}$) radicals.

1-4C-Alkoxy carbonyl-1-4C-alkyl represents one of the abovementioned 1-4C-alkyl radicals which is substituted by one of the abovementioned 1-4C-alkoxy carbonyl radicals. An example which may be mentioned is the ethoxycarbonylmethyl radical ($\text{CH}_3\text{CH}_2\text{OC(O)CH}_2\text{-}$).

Di-1-4C-alkylamino represents an amino radical which is substituted by two of the abovementioned 1-4C-alkyl radicals, which may be identical or different. Examples which may be mentioned are the dimethylamino, the diethylamino and the diisopropylamino radicals.

Di-1-4C-alkylamino-1-4C-alkyl represents one of the abovementioned 1-4C-alkyl radicals which is substituted by one of the abovementioned di-1-4C-alkylamino radicals. Examples which may be mentioned are the dimethylaminomethyl, the dimethylaminoethyl and the diethylaminoethyl radicals.

Ar-1-4C-alkyl represents one of the abovementioned 1-4C-alkyl radicals which is substituted by Ar, where Ar is as defined above. Examples which may be mentioned are the phenethyl and the benzyl radicals.

1-4C-Alkoxy-carbonylamino represents an amino radical which is substituted by one of the abovementioned 1-4C-alkoxy-carbonyl radicals. Examples which may be mentioned are the ethoxycarbonylamino and the methoxycarbonylamino radicals.

1-4C-Alkoxy-1-4C-alkoxy-carbonyl represents a carbonyl group to which is attached one of the abovementioned 1-4C-alkoxy-1-4C-alkoxy radicals. Examples which may be mentioned are the 2-(methoxy)ethoxycarbonyl ($\text{CH}_3\text{-O-CH}_2\text{CH}_2\text{-O-CO-}$) and the 2-(ethoxy)ethoxycarbonyl ($\text{CH}_3\text{CH}_2\text{-O-CH}_2\text{CH}_2\text{-O-CO-}$) radicals.

1-4C-Alkoxy-1-4C-alkoxy-carbonylamino represents an amino radical which is substituted by one of the abovementioned 1-4C-alkoxy-1-4C-alkoxy-carbonyl radicals. Examples which may be mentioned are the 2-(methoxy)-ethoxycarbonylamino and the 2-(ethoxy)ethoxycarbonylamino radicals.

1-7C-Alkylene represents straight-chain or branched 1-7C-alkylene radicals, for example the methylene ($-\text{CH}_2-$), ethylene ($-\text{CH}_2\text{-CH}_2-$), trimethylene ($-\text{CH}_2\text{-CH}_2\text{-CH}_2-$), tetramethylene ($-\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2-$), 1,2-dimethylethylene [$-\text{CH}(\text{CH}_3)\text{-CH}(\text{CH}_3)-$], 1,1-dimethylethylene [$-\text{C}(\text{CH}_3)_2\text{-CH}_2-$], 2,2-dimethylethylene [$-\text{CH}_2\text{-C}(\text{CH}_3)_2-$], isopropylidene [$-\text{C}(\text{CH}_3)_2-$], 1-methylethylene [$-\text{CH}(\text{CH}_3)\text{-CH}_2-$], pentamethylene ($-\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2-$), hexamethylene ($-\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2-$) and the heptamethylene ($-\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2-$) radicals.

3-4C-Alkenylene represents straight-chain 3-4C-alkenylene radicals, for example the 1-propenylene, the 2-propenylene, the 2-butenylene and the 3-butenylene radicals.

Suitable salts of compounds of the formula I are - depending on the substitution - in particular all acid addition salts. Particular mention may be made of the pharmacologically acceptable salts of the inorganic and organic acids customarily used in pharmacy. Those suitable are water-soluble and water-insoluble acid addition salts with acids such as, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid, sulfuric acid, acetic acid, citric acid, D-gluconic acid, benzoic acid, 2-(4-hydroxybenzoyl)benzoic acid, butyric acid, sulfosalicylic acid, maleic acid, lauric acid, malic acid, fumaric acid, succinic acid, oxalic acid, tartaric acid, embonic acid, stearic acid, toluenesulfonic acid, methanesulfonic acid or 3-hydroxy-2-naphthoic acid, where the acids are employed in the salt preparation in an equimolar ratio or a ratio differing therefrom, depending on whether the acid is a mono- or polybasic acid and on which salt is desired.

Pharmacologically unacceptable salts, which can be initially obtained, for example, as process products in the preparation of the compounds according to the invention on an industrial scale, are converted into pharmacologically acceptable salts by processes known to the person skilled in the art.

It is known to the person skilled in the art that the compounds according to the invention and their salts can, for example when they are isolated in crystalline form, comprise varying amounts of solvents. The invention therefore also embraces all solvates and, in particular, all hydrates of the compounds of the formula I, and all solvates and, in particular, all hydrates of the salts of the compounds of the formula I.

The compounds of the formula I have at least three chiral centers. The invention provides all eight feasible stereoisomers in any mixing ratio, including the pure enantiomers which are the preferred subject matter of the invention.

An exemplary preferred radical R1 is the methyl radical.

Exemplary preferred radicals R2 are the methyl and the hydroxymethyl radicals.

In the context of the present invention, R3 is preferably hydrogen.

Exemplary radicals R9 or R10 which may be mentioned are: methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, difluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxy-ethyl, 3-hydroxypropyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, carboxymethyl, carboxyethyl, carboxypropyl, methoxycarbonylmethyl, dimethylaminoethyl, diethylaminoethyl, phenyl, benzyl, 4-chlorophenyl, 4-aminophenyl, 4-chlorobenzyl, 4-difluoromethoxyphenyl, 4-trifluoromethoxyphenyl, 4-methylbenzyl, 3-methylbenzyl, 2,4-diaminophenyl, 2-methyl-4-tert-butylphenyl, 2-nitro-4-acetylphenyl, 4-fluorobenzyl, 4-nitrophenyl, 3-nitrophenyl, 3-aminophenyl, 2-methoxycarbonylamino-6-methylphenyl, 2-methoxyethoxycarbonylamino-6-methylphenyl, 2-methoxycarbonylamino-6-methylbenzyl and 2-methoxyethoxycarbonylamino-6-methylbenzyl.

Exemplary alkylene and alkenylene groups alk1 and alk2, respectively, which may be mentioned are: 1-methylethylene, 2-methylethylene, 1-phenylethylene, 2-phenylethylene, 1-propylpropylene, 3-propylpropylene, 2-aminopropylene, 2-tert-butyloxycarbonylamino-3-propylene, 2-hydroxypropylene, 2-oxopropylene, 2-carboxypropylene, 1-acetyl-1,2-dimethylethylene, 2-acetyl-1,2-dimethylethylene, 1,1-dimethyl-2-oxoethylene, 1-oxo-2,2-dimethylethylene, 1,3-dioxobutylene, 2,4-dioxobutylene, 1,2-dioxopropylene, 2,3-dioxopropylene, prop-1-enylene, prop-2-enylene, but-1-enylene, but-2-enylene, but-3-enylene, but-4-enylene, buta-1,3-dienylene, buta-2,4-dienylene, 1-oxo-but-2-enylene, 4-oxo-but-2-enylene, 1-oxo-2,2-difluoroethylene, 2-oxo-1,1-difluoroethylene, 1-oxopropylene, 3-oxopropylene, 1-carboxyethylene and 2-carboxyethylene.

Compounds which may be mentioned are those of the formula I in which

- R1 is 1-4C-alkyl,
- R2 is 1-4C-alkyl or hydroxy-1-4C-alkyl,
- R3 is hydrogen,

one of the substituents R4a and R4b is hydrogen and the other is hydrogen, hydroxyl, 1-4C-alkoxy, 1-4C-alkoxy-1-4C-alkoxy or the radical R4',

where

R4' is -O-(CH₂)_m-S(O)_n-R9,
 -S(O)_n-(CH₂)_m-OH,
 -S(O)_n-(CH₂)_m-O-R9,
 -S(O)_n-(CH₂)_m-S(O)_p-R9 or
 -S(O)_n-R9,

in which

R9 is 1-7C-alkyl, halo-1-4C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,
 m is an integer from 2 to 7,
 n is the number 0, 1 or 2 and
 p is the number 0, 1 or 2,

one of the substituents R5a and R5b is hydrogen and the other is hydrogen, hydroxyl, 1-4C-alkoxy, 1-4C-alkoxy-1-4C-alkoxy or the radical R5',

where

R5' is -O-(CH₂)_q-S(O)_r-R10,
 -S(O)_r-(CH₂)_q-OH,
 -S(O)_r-(CH₂)_q-O-R10,
 -S(O)_r-(CH₂)_q-S(O)_t-R10 or
 -S(O)_r-R10,

in which

R10 is 1-7C-alkyl, halo-1-4C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,
 q is an integer from 2 to 7,
 r is the number 0, 1 or 2 and
 t is the number 0, 1 or 2,

R6 is hydrogen, halogen or trifluoromethyl,


R7 is hydrogen or halogen and

R8 is hydrogen,

and their salts,

where one of the substituents R4a and R4b has to be R4' and/or one of the substituents R5a and R5b has to be R5'.

Compounds according to the invention which are to be emphasized are those of the formula I*



in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl or hydroxy-1-4C-alkyl,

R3 is hydrogen,

one of the substituents R4a and R4b is hydrogen and the other is hydrogen, hydroxyl, 1-4C-alkoxy, 1-4C-alkoxy-1-4C-alkoxy or the radical R4',

where

R4' is -O-(CH₂)_m-S(O)_n-R9,
 -S(O)_n-(CH₂)_m-OH,
 -S(O)_n-(CH₂)_m-O-R9,
 -S(O)_n-(CH₂)_m-S(O)_p-R9 or
 -S(O)_n-R9,

in which

R9 is 1-7C-alkyl, halo-1-4C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

m is an integer from 2 to 7,

n is the number 0, 1 or 2 and

p is the number 0, 1 or 2,

one of the substituents R5a and R5b is hydrogen and the other is hydrogen, hydroxyl, 1-4C-alkoxy, 1-4C-alkoxy-1-4C-alkoxy or the radical R5',

where

R5' is -O-(CH₂)_q-S(O)_r-R10,
 -S(O)_r-(CH₂)_q-OH,
 -S(O)_r-(CH₂)_q-O-R10,
 -S(O)_r-(CH₂)_q-S(O)_t-R10 or
 -S(O)_r-R10,

in which

R10 is 1-7C-alkyl, halo-1-4C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

q is an integer from 2 to 7,

r is the number 0, 1 or 2 and

t is the number 0, 1 or 2,

R6 is hydrogen, halogen or trifluoromethyl,

R7 is hydrogen or halogen and

R8 is hydrogen,

and their salts,

where one of the substituents R4a and R4b has to be R4' and/or one of the substituents R5a and R5b has to be R5'.

One embodiment (embodiment a) of the compounds of the formula I* to be emphasized are those in which

one of the substituents R4a and R4b is hydrogen and the other is the radical R4' and one of the substituents R5a and R5b is hydrogen and the other is hydroxyl.

A further embodiment (embodiment b) of the compounds of the formula I* to be emphasized are those in which

one of the substituents R4a and R4b is hydrogen and the other is the radical R4' and one of the substituents R5a and R5b is hydrogen and the other is 1-4C-alkoxy or 1-4C-alkoxy-1-4C-alkoxy.

A further embodiment (embodiment c) of the compounds of the formula I* to be emphasized are those in which

one of the substituents R4a and R4b is hydrogen and the other is the radical R4' and one of the substituents R5a and R5b is hydrogen and the other is the radical R5'.

A further embodiment (embodiment d) of the compounds of the formula I* to be emphasized are those in which

one of the substituents R4a and R4b is hydrogen and the other is 1-4C-alkoxy or 1-4C-alkoxy-1-4C-alkoxy and one of the substituents R5a and R5b is hydrogen and the other is the radical R5'.

Compounds I* of the embodiments a, b, c and d which are to be emphasized particularly are those in which

R1 is 1-4C-alkyl,


R2 is 1-4C-alkyl,

R3 is hydrogen,

R4' is -O-(CH₂)_m-S(O)_n-R9,




-S(O)_n-R9,

-S(O)_n-(CH₂)_m-OH or



in which R₉ is 1-4C-alkyl,

R_{5'} is $-\text{O}-(\text{CH}_2)_q-\text{S(O)}_r-\text{R}_{10}$,



in which R₁₀ is 1-4C-alkyl,

R₆ is hydrogen,

R₇ is hydrogen and

R₈ is hydrogen,

and their salts.

Preferred compounds I* of the embodiment a are those in which

R₁ is 1-4C-alkyl,




R₂ is 1-4C-alkyl,

R₃ is hydrogen,

one of the substituents R_{4a} and R_{4b} is hydrogen and the other is R_{4'},

where

R_{4'} is $-\text{O}-(\text{CH}_2)_m-\text{S(O)}_n-\text{R}_9$,



in which R₉ is 1-4C-alkyl,

m is the number 2 or 3 and

n is the number 0, 1 or 2,

one of the substituents R_{5a} and R_{5b} is hydrogen and the other is hydroxyl,

R₆ is hydrogen,

R₇ is hydrogen and

R₈ is hydrogen,

and their salts.

Preferred compounds I* of the embodiment b are those in which

R₁ is 1-4C-alkyl,


R₂ is 1-4C-alkyl,

R₃ is hydrogen,

one of the substituents R_{4a} and R_{4b} is hydrogen and the other is R_{4'},

where

R_{4'} is $-\text{O}-(\text{CH}_2)_m-\text{S(O)}_n-\text{R}_9$,



$-\text{S}(\text{O})_n-(\text{CH}_2)_m-\text{OH}$ or

$-\text{S}(\text{O})_n-(\text{CH}_2)_m-\text{O}-\text{R}_9$,

in which R_9 is 1-4C-alkyl,

m is the number 2 or 3 and

n is the number 0, 1 or 2,

one of the substituents R_{5a} and R_{5b} is hydrogen and the other is 1-4C-alkoxy or 1-4C-alkoxy-1-4C-alkoxy,

R_6 is hydrogen,

R_7 is hydrogen and

R_8 is hydrogen,

and their salts.

Preferred compounds 1^* of the embodiment c are those in which

R_1 is 1-4C-alkyl,

R_2 is 1-4C-alkyl,

R_3 is hydrogen,

one of the substituents R_{4a} and R_{4b} is hydrogen and the other is $\text{R}_{4'}$,

where

$\text{R}_{4'}$ is $-\text{O}-(\text{CH}_2)_m-\text{S}(\text{O})_n-\text{R}_9$,

$-\text{S}(\text{O})_n-\text{R}_9$,

$-\text{S}(\text{O})_n-(\text{CH}_2)_m-\text{OH}$ or

$-\text{S}(\text{O})_n-(\text{CH}_2)_m-\text{O}-\text{R}_9$,

in which R_9 is 1-4C-alkyl,

m is the number 2 or 3 and

n is the number 0, 1 or 2,

one of the substituents R_{5a} and R_{5b} is hydrogen and the other is $\text{R}_{5'}$,

where

$\text{R}_{5'}$ is $-\text{O}-(\text{CH}_2)_q-\text{S}(\text{O})_r-\text{R}_{10}$,

$-\text{S}(\text{O})_r-\text{R}_{10}$,

$-\text{S}(\text{O})_r-(\text{CH}_2)_q-\text{OH}$ or

$-\text{S}(\text{O})_r-(\text{CH}_2)_q-\text{O}-\text{R}_{10}$,

in which R_{10} is 1-4C-alkyl,

q is the number 2 or 3 and

r is the number 0, 1 or 2,

R_6 is hydrogen,

R_7 is hydrogen and

R_8 is hydrogen,

and their salts.

Preferred compounds I* of the embodiment d are those in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl,

R3 is hydrogen,

one of the substituents R4a and R4b is hydrogen and the other is 1-4C-alkoxy or 1-4C-alkoxy-1-4C-alkoxy,

one of the substituents R5a and R5b is hydrogen and the other is R5',

where

R5' is -O-(CH₂)_q-S(O)_r-R10,

-S(O)_r-R10,

-S(O)_r-(CH₂)_q-OH or

-S(O)_r-(CH₂)_q-O-R10,

in which R10 is 1-4C-alkyl,

q is the number 2 or 3 and

r is the number 0, 1 or 2,

R6 is hydrogen,

R7 is hydrogen and


R8 is hydrogen,

and their salts.

Selected preferred compounds I* are those of the embodiments a, b, c and d in which R5b is hydrogen.

Selected particularly preferred compounds I* are those of the embodiments a, b, c and d in which R5b is hydrogen and R4a is hydrogen.

The following preferred compounds according to the invention selected by way of example may be mentioned specifically using formula I** below with the substituent definitions for R4a, R4b and R5a of Table 1 (Tab. 1) below:



Tab. 1

R4a	R4b	R5a
H	-OCH ₂ CH ₂ SCH ₃	-OH
H	-OCH ₂ CH ₂ SOCH ₃	-OH
H	-OCH ₂ CH ₂ SO ₂ CH ₃	-OH
H	-SCH ₂ CH ₂ OCH ₃	-OH
H	-SOCH ₂ CH ₂ OCH ₃	-OH
H	-SO ₂ CH ₂ CH ₂ OCH ₃	-OH
H	-SCH ₂ CH ₂ OH	-OH
H	-SCH ₃	-OH
H	-SOCH ₃	-OH
H	-SO ₂ CH ₃	-OH
H	-SCH ₂ CH ₃	-OH
H	-SOCH ₂ CH ₃	-OH
H	-SO ₂ CH ₂ CH ₃	-OH
H	-SCH ₂ CH ₂ CH ₃	-OH
H	-SOCH ₂ CH ₂ CH ₃	-OH
H	-SO ₂ CH ₂ CH ₂ CH ₃	-OH
H	-SCH(CH ₃) ₂	-OH
H	-SOCH(CH ₃) ₂	-OH
H	-SO ₂ CH(CH ₃) ₂	-OH
H	-SCH ₂ COOCH ₂ CH ₃	-OH
H	-SOCH ₂ COOCH ₂ CH ₃	-OH
H	-SO ₂ CH ₂ COOCH ₂ CH ₃	-OH
-OCH ₂ CH ₂ SCH ₃	H	-OH
-OCH ₂ CH ₂ SOCH ₃	H	-OH
-OCH ₂ CH ₂ SO ₂ CH ₃	H	-OH
-SCH ₂ CH ₂ OCH ₃	H	-OH
-SOCH ₂ CH ₂ OCH ₃	H	-OH
-SO ₂ CH ₂ CH ₂ OCH ₃	H	-OH
-SCH ₂ CH ₂ OH	H	-OH
-SCH ₃	H	-OH
-SOCH ₃	H	-OH
-SO ₂ CH ₃	H	-OH
-SCH ₂ CH ₃	H	-OH
-SOCH ₂ CH ₃	H	-OH
-SO ₂ CH ₂ CH ₃	H	-OH

R4a	R4b	R5a
-SCH ₂ CH ₂ CH ₃	H	-OH
-SOCH ₂ CH ₂ CH ₃	H	-OH
-SO ₂ CH ₂ CH ₂ CH ₃	H	-OH
-SCH(CH ₃) ₂	H	-OH
-SOCH(CH ₃) ₂	H	-OH
-SO ₂ CH(CH ₃) ₂	H	-OH
-SCH ₂ COOCH ₂ CH ₃	H	-OH
-SOCH ₂ COOCH ₂ CH ₃	H	-OH
-SO ₂ CH ₂ COOCH ₂ CH ₃	H	-OH
H	-OCH ₂ CH ₂ SCH ₃	-OCH ₃
H	-OCH ₂ CH ₂ SOCH ₃	-OCH ₃
H	-OCH ₂ CH ₂ SO ₂ CH ₃	-OCH ₃
H	-SCH ₂ CH ₂ OCH ₃	-OCH ₃
H	-SOCH ₂ CH ₂ OCH ₃	-OCH ₃
H	-SO ₂ CH ₂ CH ₂ OCH ₃	-OCH ₃
H	-SCH ₂ CH ₂ OH	-OCH ₃
H	-SCH ₃	-OCH ₃
H	-SOCH ₃	-OCH ₃
H	-SO ₂ CH ₃	-OCH ₃
H	-SCH ₂ CH ₃	-OCH ₃
H	-SOCH ₂ CH ₃	-OCH ₃
H	-SO ₂ CH ₂ CH ₃	-OCH ₃
H	-SCH ₂ CH ₂ CH ₃	-OCH ₃
H	-SOCH ₂ CH ₂ CH ₃	-OCH ₃
H	-SO ₂ CH ₂ CH ₂ CH ₃	-OCH ₃
H	-SCH(CH ₃) ₂	-OCH ₃
H	-SOCH(CH ₃) ₂	-OCH ₃
H	-SO ₂ CH(CH ₃) ₂	-OCH ₃
H	-SCH ₂ COOCH ₂ CH ₃	-OCH ₃
H	-SOCH ₂ COOCH ₂ CH ₃	-OCH ₃
H	-SO ₂ CH ₂ COOCH ₂ CH ₃	-OCH ₃
H	-OCH ₂ CH ₂ SCH ₃	-OCH ₂ CH ₂ OCH ₃
H	-OCH ₂ CH ₂ SOCH ₃	-OCH ₂ CH ₂ OCH ₃
H	-OCH ₂ CH ₂ SO ₂ CH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SCH ₂ CH ₂ OCH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SOCH ₂ CH ₂ OCH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SO ₂ CH ₂ CH ₂ OCH ₃	-OCH ₂ CH ₂ OCH ₃

R4a	R4b	R5a
H	-SCH ₂ CH ₂ OH	-OCH ₂ CH ₂ OCH ₃
H	-SCH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SOCH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SO ₂ CH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SCH ₂ CH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SOCH ₂ CH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SO ₂ CH ₂ CH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SCH ₂ CH ₂ CH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SOCH ₂ CH ₂ CH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SO ₂ CH ₂ CH ₂ CH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SCH(CH ₃) ₂	-OCH ₂ CH ₂ OCH ₃
H	-SOCH(CH ₃) ₂	-OCH ₂ CH ₂ OCH ₃
H	-SO ₂ CH(CH ₃) ₂	-OCH ₂ CH ₂ OCH ₃
H	-SCH ₂ COOCH ₂ CH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SOCH ₂ COOCH ₂ CH ₃	-OCH ₂ CH ₂ OCH ₃
H	-SO ₂ CH ₂ COOCH ₂ CH ₃	-OCH ₂ CH ₂ OCH ₃
-OCH ₂ CH ₂ SCH ₃	H	-OCH ₃
-OCH ₂ CH ₂ SOCH ₃	H	-OCH ₃
-OCH ₂ CH ₂ SO ₂ CH ₃	H	-OCH ₃
-SCH ₂ CH ₂ OCH ₃	H	-OCH ₃
-SOCH ₂ CH ₂ OCH ₃	H	-OCH ₃
-SO ₂ CH ₂ CH ₂ OCH ₃	H	-OCH ₃
-SCH ₂ CH ₂ OH	H	-OCH ₃
-SCH ₃	H	-OCH ₃
-SOCH ₃	H	-OCH ₃
-SO ₂ CH ₃	H	-OCH ₃
-SCH ₂ CH ₃	H	-OCH ₃
-SOCH ₂ CH ₃	H	-OCH ₃
-SO ₂ CH ₂ CH ₃	H	-OCH ₃
-SCH ₂ CH ₂ CH ₃	H	-OCH ₃
-SOCH ₂ CH ₂ CH ₃	H	-OCH ₃
-SO ₂ CH ₂ CH ₂ CH ₃	H	-OCH ₃
-SCH(CH ₃) ₂	H	-OCH ₃
-SOCH(CH ₃) ₂	H	-OCH ₃
-SO ₂ CH(CH ₃) ₂	H	-OCH ₃
-SCH ₂ COOCH ₂ CH ₃	H	-OCH ₃
-SOCH ₂ COOCH ₂ CH ₃	H	-OCH ₃

R4a	R4b	R5a
-SO ₂ CH ₂ COOCH ₂ CH ₃	H	-OCH ₃
-OCH ₂ CH ₂ SCH ₃	H	-OCH ₂ CH ₂ OCH ₃
-OCH ₂ CH ₂ SOCH ₃	H	-OCH ₂ CH ₂ OCH ₃
-OCH ₂ CH ₂ SO ₂ CH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SCH ₂ CH ₂ OCH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SOCH ₂ CH ₂ OCH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SO ₂ CH ₂ CH ₂ OCH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SCH ₂ CH ₂ OH	H	-OCH ₂ CH ₂ OCH ₃
-SCH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SOCH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SO ₂ CH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SCH ₂ CH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SOCH ₂ CH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SO ₂ CH ₂ CH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SCH ₂ CH ₂ CH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SOCH ₂ CH ₂ CH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SO ₂ CH ₂ CH ₂ CH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SCH(CH ₃) ₂	H	-OCH ₂ CH ₂ OCH ₃
-SOCH(CH ₃) ₂	H	-OCH ₂ CH ₂ OCH ₃
-SO ₂ CH(CH ₃) ₂	H	-OCH ₂ CH ₂ OCH ₃
-SCH ₂ COOCH ₂ CH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SOCH ₂ COOCH ₂ CH ₃	H	-OCH ₂ CH ₂ OCH ₃
-SO ₂ CH ₂ COOCH ₂ CH ₃	H	-OCH ₂ CH ₂ OCH ₃
-OCH ₂ CH ₂ SCH ₃	H	-OCH ₂ CH ₂ SCH ₃
-OCH ₂ CH ₂ SOCH ₃	H	-OCH ₂ CH ₂ SCH ₃
-OCH ₂ CH ₂ SO ₂ CH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SCH ₂ CH ₂ OCH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SOCH ₂ CH ₂ OCH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SO ₂ CH ₂ CH ₂ OCH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SCH ₂ CH ₂ OH	H	-OCH ₂ CH ₂ SCH ₃
-SCH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SOCH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SO ₂ CH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SCH ₂ CH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SOCH ₂ CH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SO ₂ CH ₂ CH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SCH ₂ CH ₂ CH ₃	H	-OCH ₂ CH ₂ SCH ₃

R4a	R4b	R5a
-SOCH ₂ CH ₂ CH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SO ₂ CH ₂ CH ₂ CH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SCH(CH ₃) ₂	H	-OCH ₂ CH ₂ SCH ₃
-SOCH(CH ₃) ₂	H	-OCH ₂ CH ₂ SCH ₃
-SO ₂ CH(CH ₃) ₂	H	-OCH ₂ CH ₂ SCH ₃
-SCH ₂ COOCH ₂ CH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SOCH ₂ COOCH ₂ CH ₃	H	-OCH ₂ CH ₂ SCH ₃
-SO ₂ CH ₂ COOCH ₂ CH ₃	H	-OCH ₂ CH ₂ SCH ₃
-OCH ₂ CH ₂ SCH ₃	H	-SCH ₂ CH ₂ OH
-OCH ₂ CH ₂ SOCH ₃	H	-SCH ₂ CH ₂ OH
-OCH ₂ CH ₂ SO ₂ CH ₃	H	-SCH ₂ CH ₂ OH
-SCH ₂ CH ₂ OCH ₃	H	-SCH ₂ CH ₂ OH
-SOCH ₂ CH ₂ OCH ₃	H	-SCH ₂ CH ₂ OH
-SO ₂ CH ₂ CH ₂ OCH ₃	H	-SCH ₂ CH ₂ OH
-SCH ₂ CH ₂ OH	H	-SCH ₂ CH ₂ OH
-SCH ₃	H	-SCH ₂ CH ₂ OH
-SOCH ₃	H	-SCH ₂ CH ₂ OH
-SO ₂ CH ₃	H	-SCH ₂ CH ₂ OH
-SCH ₂ CH ₃	H	-SCH ₂ CH ₂ OH
-SOCH ₂ CH ₃	H	-SCH ₂ CH ₂ OH
-SO ₂ CH ₂ CH ₃	H	-SCH ₂ CH ₂ OH
-SCH ₂ CH ₂ CH ₃	H	-SCH ₂ CH ₂ OH
-SOCH ₂ CH ₂ CH ₃	H	-SCH ₂ CH ₂ OH
-SO ₂ CH ₂ CH ₂ CH ₃	H	-SCH ₂ CH ₂ OH
-SCH(CH ₃) ₂	H	-SCH ₂ CH ₂ OH
-SOCH(CH ₃) ₂	H	-SCH ₂ CH ₂ OH
-SO ₂ CH(CH ₃) ₂	H	-SCH ₂ CH ₂ OH
-SCH ₂ COOCH ₂ CH ₃	H	-SCH ₂ CH ₂ OH
-SOCH ₂ COOCH ₂ CH ₃	H	-SCH ₂ CH ₂ OH
-SO ₂ CH ₂ COOCH ₂ CH ₃	H	-SCH ₂ CH ₂ OH
-OCH ₂ CH ₂ SCH ₃	H	-SCH ₃
-OCH ₂ CH ₂ SOCH ₃	H	-SCH ₃
-OCH ₂ CH ₂ SO ₂ CH ₃	H	-SCH ₃
-SCH ₂ CH ₂ OCH ₃	H	-SCH ₃
-SOCH ₂ CH ₂ OCH ₃	H	-SCH ₃
-SO ₂ CH ₂ CH ₂ OCH ₃	H	-SCH ₃
-SCH ₂ CH ₂ OH	H	-SCH ₃

R4a	R4b	R5a
-SCH ₃	H	-SCH ₃
-SOCH ₃	H	-SCH ₃
-SO ₂ CH ₃	H	-SCH ₃
-SCH ₂ CH ₃	H	-SCH ₃
-SOCH ₂ CH ₃	H	-SCH ₃
-SO ₂ CH ₂ CH ₃	H	-SCH ₃
-SCH ₂ CH ₂ CH ₃	H	-SCH ₃
-SOCH ₂ CH ₂ CH ₃	H	-SCH ₃
-SO ₂ CH ₂ CH ₂ CH ₃	H	-SCH ₃
-SCH(CH ₃) ₂	H	-SCH ₃
-SOCH(CH ₃) ₂	H	-SCH ₃
-SO ₂ CH(CH ₃) ₂	H	-SCH ₃
-SCH ₂ COOCH ₂ CH ₃	H	-SCH ₃
-SOCH ₂ COOCH ₂ CH ₃	H	-SCH ₃
-SO ₂ CH ₂ COOCH ₂ CH ₃	H	-SCH ₃
-OCH ₂ CH ₂ SCH ₃	H	-SCH ₂ CH ₂ OCH ₃
-OCH ₂ CH ₂ SOCH ₃	H	-SCH ₂ CH ₂ OCH ₃
-OCH ₂ CH ₂ SO ₂ CH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SCH ₂ CH ₂ OCH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SOCH ₂ CH ₂ OCH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SO ₂ CH ₂ CH ₂ OCH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SCH ₂ CH ₂ OH	H	-SCH ₂ CH ₂ OCH ₃
-SCH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SOCH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SO ₂ CH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SCH ₂ CH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SOCH ₂ CH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SO ₂ CH ₂ CH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SCH ₂ CH ₂ CH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SOCH ₂ CH ₂ CH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SO ₂ CH ₂ CH ₂ CH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SCH(CH ₃) ₂	H	-SCH ₂ CH ₂ OCH ₃
-SOCH(CH ₃) ₂	H	-SCH ₂ CH ₂ OCH ₃
-SO ₂ CH(CH ₃) ₂	H	-SCH ₂ CH ₂ OCH ₃
-SCH ₂ COOCH ₂ CH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SOCH ₂ COOCH ₂ CH ₃	H	-SCH ₂ CH ₂ OCH ₃
-SO ₂ CH ₂ COOCH ₂ CH ₃	H	-SCH ₂ CH ₂ OCH ₃

R4a	R4b	R5a
H	-OCH ₂ CH ₂ SCH ₃	-OCH ₂ CH ₂ SCH ₃
H	-OCH ₂ CH ₂ SOCH ₃	-OCH ₂ CH ₂ SCH ₃
H	-OCH ₂ CH ₂ SO ₂ CH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SCH ₂ CH ₂ OCH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SOCH ₂ CH ₂ OCH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SO ₂ CH ₂ CH ₂ OCH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SCH ₂ CH ₂ OH	-OCH ₂ CH ₂ SCH ₃
H	-SCH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SOCH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SO ₂ CH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SCH ₂ CH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SOCH ₂ CH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SO ₂ CH ₂ CH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SCH ₂ CH ₂ CH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SOCH ₂ CH ₂ CH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SO ₂ CH ₂ CH ₂ CH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SCH(CH ₃) ₂	-OCH ₂ CH ₂ SCH ₃
H	-SOCH(CH ₃) ₂	-OCH ₂ CH ₂ SCH ₃
H	-SO ₂ CH(CH ₃) ₂	-OCH ₂ CH ₂ SCH ₃
H	-SCH ₂ COOCH ₂ CH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SOCH ₂ COOCH ₂ CH ₃	-OCH ₂ CH ₂ SCH ₃
H	-SO ₂ CH ₂ COOCH ₂ CH ₃	-OCH ₂ CH ₂ SCH ₃
H	-OCH ₂ CH ₂ SCH ₃	-SCH ₂ CH ₂ OH
H	-OCH ₂ CH ₂ SOCH ₃	-SCH ₂ CH ₂ OH
H	-OCH ₂ CH ₂ SO ₂ CH ₃	-SCH ₂ CH ₂ OH
H	-SCH ₂ CH ₂ OCH ₃	-SCH ₂ CH ₂ OH
H	-SOCH ₂ CH ₂ OCH ₃	-SCH ₂ CH ₂ OH
H	-SO ₂ CH ₂ CH ₂ OCH ₃	-SCH ₂ CH ₂ OH
H	-SCH ₂ CH ₂ OH	-SCH ₂ CH ₂ OH
H	-SCH ₃	-SCH ₂ CH ₂ OH
H	-SOCH ₃	-SCH ₂ CH ₂ OH
H	-SO ₂ CH ₃	-SCH ₂ CH ₂ OH
H	-SCH ₂ CH ₃	-SCH ₂ CH ₂ OH
H	-SOCH ₂ CH ₃	-SCH ₂ CH ₂ OH
H	-SO ₂ CH ₂ CH ₃	-SCH ₂ CH ₂ OH
H	-SCH ₂ CH ₂ CH ₃	-SCH ₂ CH ₂ OH
H	-SOCH ₂ CH ₂ CH ₃	-SCH ₂ CH ₂ OH

R4a	R4b	R5a
H	-SO ₂ CH ₂ CH ₂ CH ₃	-SCH ₂ CH ₂ OH
H	-SCH(CH ₃) ₂	-SCH ₂ CH ₂ OH
H	-SOCH(CH ₃) ₂	-SCH ₂ CH ₂ OH
H	-SO ₂ CH(CH ₃) ₂	-SCH ₂ CH ₂ OH
H	-SCH ₂ COOCH ₂ CH ₃	-SCH ₂ CH ₂ OH
H	-SOCH ₂ COOCH ₂ CH ₃	-SCH ₂ CH ₂ OH
H	-SO ₂ CH ₂ COOCH ₂ CH ₃	-SCH ₂ CH ₂ OH
H	-OCH ₂ CH ₂ SCH ₃	-SCH ₃
H	-OCH ₂ CH ₂ SOCH ₃	-SCH ₃
H	-OCH ₂ CH ₂ SO ₂ CH ₃	-SCH ₃
H	-SCH ₂ CH ₂ OCH ₃	-SCH ₃
H	-SOCH ₂ CH ₂ OCH ₃	-SCH ₃
H	-SO ₂ CH ₂ CH ₂ OCH ₃	-SCH ₃
H	-SCH ₂ CH ₂ OH	-SCH ₃
H	-SCH ₃	-SCH ₃
H	-SOCH ₃	-SCH ₃
H	-SO ₂ CH ₃	-SCH ₃
H	-SCH ₂ CH ₃	-SCH ₃
H	-SOCH ₂ CH ₃	-SCH ₃
H	-SO ₂ CH ₂ CH ₃	-SCH ₃
H	-SCH ₂ CH ₂ CH ₃	-SCH ₃
H	-SOCH ₂ CH ₂ CH ₃	-SCH ₃
H	-SO ₂ CH ₂ CH ₂ CH ₃	-SCH ₃
H	-SCH(CH ₃) ₂	-SCH ₃
H	-SOCH(CH ₃) ₂	-SCH ₃
H	-SO ₂ CH(CH ₃) ₂	-SCH ₃
H	-SCH ₂ COOCH ₂ CH ₃	-SCH ₃
H	-SOCH ₂ COOCH ₂ CH ₃	-SCH ₃
H	-SO ₂ CH ₂ COOCH ₂ CH ₃	-SCH ₃
H	-OCH ₂ CH ₂ SCH ₃	-SCH ₂ CH ₂ OCH ₃
H	-OCH ₂ CH ₂ SOCH ₃	-SCH ₂ CH ₂ OCH ₃
H	-OCH ₂ CH ₂ SO ₂ CH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SCH ₂ CH ₂ OCH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SOCH ₂ CH ₂ OCH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SO ₂ CH ₂ CH ₂ OCH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SCH ₂ CH ₂ OH	-SCH ₂ CH ₂ OCH ₃
H	-SCH ₃	-SCH ₂ CH ₂ OCH ₃

R4a	R4b	R5a
H	-SOCH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SO ₂ CH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SCH ₂ CH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SOCH ₂ CH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SO ₂ CH ₂ CH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SCH ₂ CH ₂ CH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SOCH ₂ CH ₂ CH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SO ₂ CH ₂ CH ₂ CH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SCH(CH ₃) ₂	-SCH ₂ CH ₂ OCH ₃
H	-SOCH(CH ₃) ₂	-SCH ₂ CH ₂ OCH ₃
H	-SO ₂ CH(CH ₃) ₂	-SCH ₂ CH ₂ OCH ₃
H	-SCH ₂ COOCH ₂ CH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SOCH ₂ COOCH ₂ CH ₃	-SCH ₂ CH ₂ OCH ₃
H	-SO ₂ CH ₂ COOCH ₂ CH ₃	-SCH ₂ CH ₂ OCH ₃
H	-OCH ₃	-OCH ₂ CH ₂ SCH ₃
H	-OCH ₃	-OCH ₂ CH ₂ SOCH ₃
H	-OCH ₃	-OCH ₂ CH ₂ SO ₂ CH ₃
H	-OCH ₃	-SCH ₂ CH ₂ OCH ₃
H	-OCH ₃	-SOCH ₂ CH ₂ OCH ₃
H	-OCH ₃	-SO ₂ CH ₂ CH ₂ OCH ₃
H	-OCH ₃	-SCH ₂ CH ₂ OH
H	-OCH ₃	-SCH ₃
H	-OCH ₃	-SOCH ₃
H	-OCH ₃	-SO ₂ CH ₃
H	-OCH ₃	-SCH ₂ CH ₃
H	-OCH ₃	-SOCH ₂ CH ₃
H	-OCH ₃	-SO ₂ CH ₂ CH ₃
H	-OCH ₃	-SCH ₂ CH ₂ CH ₃
H	-OCH ₃	-SOCH ₂ CH ₂ CH ₃
H	-OCH ₃	-SO ₂ CH ₂ CH ₂ CH ₃
H	-OCH ₃	-SCH(CH ₃) ₂
H	-OCH ₃	-SOCH(CH ₃) ₂
H	-OCH ₃	-SO ₂ CH(CH ₃) ₂
H	-OCH ₃	-SCH ₂ COOCH ₂ CH ₃
H	-OCH ₃	-SOCH ₂ COOCH ₂ CH ₃
H	-OCH ₃	-SO ₂ CH ₂ COOCH ₂ CH ₃
H	-OCH ₃	-OCH ₂ CH ₂ OCH ₃

R4a	R4b	R5a
H	-OCH ₂ CH ₂ OCH ₃	-OCH ₂ CH ₂ SOCH ₃
H	-OCH ₂ CH ₂ OCH ₃	-OCH ₂ CH ₂ SO ₂ CH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SCH ₂ CH ₂ OCH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SOCH ₂ CH ₂ OCH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SO ₂ CH ₂ CH ₂ OCH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SCH ₂ CH ₂ OH
H	-OCH ₂ CH ₂ OCH ₃	-SCH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SOCH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SO ₂ CH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SCH ₂ CH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SOCH ₂ CH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SO ₂ CH ₂ CH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SCH ₂ CH ₂ CH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SOCH ₂ CH ₂ CH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SO ₂ CH ₂ CH ₂ CH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SCH(CH ₃) ₂
H	-OCH ₂ CH ₂ OCH ₃	-SOCH(CH ₃) ₂
H	-OCH ₂ CH ₂ OCH ₃	-SO ₂ CH(CH ₃) ₂
H	-OCH ₂ CH ₂ OCH ₃	-SCH ₂ COOCH ₂ CH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SOCH ₂ COOCH ₂ CH ₃
H	-OCH ₂ CH ₂ OCH ₃	-SO ₂ CH ₂ COOCH ₂ CH ₃
-OCH ₃	H	-OCH ₂ CH ₂ SCH ₃
-OCH ₃	H	-OCH ₂ CH ₂ SOCH ₃
-OCH ₃	H	-OCH ₂ CH ₂ SO ₂ CH ₃
-OCH ₃	H	-SCH ₂ CH ₂ OCH ₃
-OCH ₃	H	-SOCH ₂ CH ₂ OCH ₃
-OCH ₃	H	-SO ₂ CH ₂ CH ₂ OCH ₃
-OCH ₃	H	-SCH ₂ CH ₂ OH
-OCH ₃	H	-SCH ₃
-OCH ₃	H	-SOCH ₃
-OCH ₃	H	-SO ₂ CH ₃
-OCH ₃	H	-SCH ₂ CH ₃
-OCH ₃	H	-SOCH ₂ CH ₃
-OCH ₃	H	-SO ₂ CH ₂ CH ₃
-OCH ₃	H	-SCH ₂ CH ₂ CH ₃
-OCH ₃	H	-SOCH ₂ CH ₂ CH ₃
-OCH ₃	H	-SO ₂ CH ₂ CH ₂ CH ₃

R4a	R4b	R5a
-OCH ₃	H	-SCH(CH ₃) ₂
-OCH ₃	H	-SOCH(CH ₃) ₂
-OCH ₃	H	-SO ₂ CH(CH ₃) ₂
-OCH ₃	H	-SCH ₂ COOCH ₂ CH ₃
-OCH ₃	H	-SOCH ₂ COOCH ₂ CH ₃
-OCH ₃	H	-SO ₂ CH ₂ COOCH ₂ CH ₃
-OCH ₂ CH ₂ OCH ₃	H	-OCH ₂ CH ₂ SCH ₃
-OCH ₂ CH ₂ OCH ₃	H	-OCH ₂ CH ₂ SOCH ₃
-OCH ₂ CH ₂ OCH ₃	H	-OCH ₂ CH ₂ SO ₂ CH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SCH ₂ CH ₂ OCH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SOCH ₂ CH ₂ OCH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SO ₂ CH ₂ CH ₂ OCH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SCH ₂ CH ₂ OH
-OCH ₂ CH ₂ OCH ₃	H	-SCH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SOCH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SO ₂ CH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SCH ₂ CH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SOCH ₂ CH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SO ₂ CH ₂ CH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SCH ₂ CH ₂ CH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SOCH ₂ CH ₂ CH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SO ₂ CH ₂ CH ₂ CH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SCH(CH ₃) ₂
-OCH ₂ CH ₂ OCH ₃	H	-SOCH(CH ₃) ₂
-OCH ₂ CH ₂ OCH ₃	H	-SO ₂ CH(CH ₃) ₂
-OCH ₂ CH ₂ OCH ₃	H	-SCH ₂ COOCH ₂ CH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SOCH ₂ COOCH ₂ CH ₃
-OCH ₂ CH ₂ OCH ₃	H	-SO ₂ CH ₂ COOCH ₂ CH ₃

The compounds according to the invention are prepared, for example, starting from the compounds of the formula I known from WO98/42707 in which at least one of the substituents R4a, R4b, R5a and R5b is hydroxyl (referred to as "starting materials" hereinbelow). Starting with the starting materials, the compounds according to the invention can be prepared by various routes, depending on which end product is desired, for example as outlined below:

- a) by acid-catalyzed etherification of the starting materials with compounds of the formula R4'-H or R5'-H.

- b) by reacting the starting materials (if appropriate after they have been sulfidized) with compounds of the formula $R9-S(O)_n-(CH_2)_m-Hal$, $HO-(CH_2)_m-Hal$, $R9-O-(CH_2)_m-Hal$, $R9-S(O)_p-(CH_2)_m-Hal$, $R9-Hal$, $R9-S(O)_n-alk1-Hal$, $HO-alk1-Hal$, $R9-O-alk1-Hal$ or $R9-S(O)_p-alk1-Hal$ or $R10-S(O)_r-(CH_2)_q-Hal$, $HO-(CH_2)_q-Hal$, $R10-O-(CH_2)_q-Hal$, $R10-S(O)_t-(CH_2)_q-Hal$, $R10-Hal$, $R10-S(O)_r-alk2-Hal$, $HO-alk2-Hal$, $R10-O-alk2-Hal$ or $R10-S(O)_t-alk2-Hal$, where Hal is a halogen atom, preferably a chlorine, bromine or iodine atom, preferably in the presence of an auxiliary base or after prior deprotonation of the starting material.

Depending on the nature of the radical $R4'$ or $R5'$, it may be advantageous not to introduce the entire radical $R4'$ or $R5'$ into the starting materials by the appropriate reaction. Rather, it is also feasible to employ a two- or multi-step procedure, as described in an exemplary manner in the European Patent Application 235 575, for example.

For the compounds of the formula I according to the invention in which n , p , r and/or t are different from the number 0, the reaction according to a) or b) is, if appropriate, followed by oxidation.

The oxidation of the sulfides to the sulfoxides or sulfones is carried out under the conditions known to the person skilled in the art for oxidizing sulfides to sulfoxides and sulfones respectively, [see, for example, J. Drabowicz and M. Mikołajczyk, Organic preparations and procedures int. 14(1-2), 45-89 (1982) or E. Block in S. Patai, The Chemistry of Functional Groups, Supplement E. Part 1, pp. 539-608, John Wiley and Sons (Interscience Publication), 1980]. Suitable oxidizing agents are all reagents which are customarily used for oxidizing sulfides to sulfoxides or sulfones.

Sulfoxides are optically active compounds, so that for n , p , r and/or $t = 1$, one or more further chiral centers are introduced into the molecule. With respect to the sulfoxides, the invention therefore also embraces both the enantiomers and diastereomers and their mixtures and racemates. The enantiomers can be separated in a manner known per se (for example by preparing and separating corresponding diastereoisomeric compounds).

The compounds of the formulae $R4'-H$, $R5'-H$, $R9-S(O)_n-(CH_2)_m-Hal$, $HO-(CH_2)_m-Hal$, $R9-O-(CH_2)_m-Hal$, $R9-S(O)_p-(CH_2)_m-Hal$, $R9-Hal$, $R9-S(O)_n-alk1-Hal$, $HO-alk1-Hal$, $R9-O-alk1-Hal$, $R9-S(O)_p-alk1-Hal$, $R10-S(O)_r-(CH_2)_q-Hal$, $HO-(CH_2)_q-Hal$, $R10-O-(CH_2)_q-Hal$, $R10-S(O)_t-(CH_2)_q-Hal$, $R10-Hal$, $R10-S(O)_r-alk2-Hal$, $HO-alk2-Hal$, $R10-O-alk2-Hal$, $R10-S(O)_t-alk2-Hal$ and the compounds required for their preparation are known, or they can be prepared in a manner known per se.

The following examples illustrate the invention in more detail, without limiting it. The compounds according to the invention can be prepared in a manner analogous to that described in the examples. The abbreviation RT stands for room temperature, h stands for hour(s), m.p. stands for melting point and decomp. stands for decomposition.

Examples

1. **(7R,8R,9R)-2,3-dimethyl-8-hydroxy-7-(2-methylthioethoxy)-9-phenyl-7,8,9,10-tetrahydroimidazo[1,2-h][1,7]naphthyridine**

1 g of (7R,8R,9R)-7,8-dihydroxy-2,3-dimethyl-9-phenyl-7,8,9,10-tetrahydroimidazo[1,2-h][1,7]naphthyridine, dissolved in 30 ml of dioxane and 5 ml of dimethyl-formamide, is admixed with 0.63 g of concentrated sulfuric acid and 1 g of 2-methylmercaptoethanol. The mixture is stirred at RT for 3 h, poured into ice-water (200 ml), adjusted to pH 8 using 2 N aqueous sodium hydroxide solution and extracted three times with methylene chloride. The solvent is stripped off under reduced pressure and the oily residue that remains is purified on silica gel (mobile phase: diethyl ether). This gives 80 mg of the title compound of m.p. 118-9°C (diisopropyl ether).

2. **(7S,8R,9R)-2,3-dimethyl-8-hydroxy-7-(2-methylthioethoxy)-9-phenyl-7,8,9,10-tetrahydroimidazo[1,2-h][1,7]naphthyridine**

The mother liquor from Example 1 gives, after purification on silica gel (mobile phase: diethyl ether), 40 mg of the title compound of m.p. 109-10°C.

3. **(7R,8R,9R)-2,3-dimethyl-8-hydroxy-7-(2-methylsulphinylethoxy)-9-phenyl-7,8,9,10-tetrahydroimidazo[1,2-h][1,7]naphthyridine**

130 mg of sodium metaperiodate, dissolved in 1 ml of water, is added to a solution of 200 mg of (7R,8R,9R)-2,3-dimethyl-8-hydroxy-7-(2-methylthioethoxy)-9-phenyl-7,8,9,10-tetrahydroimidazo[1,2-h][1,7]naphthyridine in 5 ml of methanol. After two hours stirring at room temperature, 50 ml of water are added. The mixture is extracted three times with dichloromethane, the organic phases are collectively washed with water and dried over sodium sulphate. The solvent is removed in vacuo and the remaining residue is purified by chromatography on silica gel (eluent: ethylacetate/methanol = 10/1). 100 mg of the title compound of m.p. 105 -106°C are obtained.

4. **(7S,8R,9R)-2,3-dimethyl-8-hydroxy-7-(2-methylsulphinylethoxy)-9-phenyl-7,8,9,10-tetrahydroimidazo[1,2-h][1,7]naphthyridine**

80 mg of the title compound of m.p. 85 - 87°C are obtained by oxidation of 200 mg of (7S,8R,9R)-2,3-dimethyl-8-hydroxy-7-(2-methylthioethoxy)-9-phenyl-7,8,9,10-tetrahydro-imidazo[1,2-h][1,7]naphthyridine analogously to Example 3.

5. (7R,8R,9R)-2,3-dimethyl-8-hydroxy-7-(ethylthio)-9-phenyl-7,8,9,10-tetrahydro-imidazo-[1,2-h][1,7]naphthyridine

1.9 g of concentrated sulphuric acid are added slowly at room temperature with stirring to 3 g of (7R,8R,9R)-7,8-dihydroxy-2,3-dimethyl-9-phenyl-7,8,9,10-tetra-hydroimidazo[1,2-h][1,7]naphthyridine, dissolved in a mixture of 40 ml of dry dioxan and 4 ml of ethylsulfide: The temperature is brought to 50°C by external heating and is then kept for 15 min. After cooling to room temperature, the mixture is poured into 300 g of a mixture of crushed ice and water. The pH is adjusted to neutral by adding aqueous 2N sodium hydroxide solution. The mixture is extracted three times (50 ml each) with ethyl acetate, the organic layers are collectively washed with water and dried over potassium carbonate. The solvent is stripped off under reduced pressure and the solid residue that remains is purified by chromatography on silica gel (mobile phase: diethyl ether). 700 mg of the title compound of m.p. 220 - 223°C are obtained after crystallisation from 2-propanol.

6. (7S,8R,9R)-2,3-dimethyl-8-hydroxy-7-(ethylthio)-9-phenyl-7,8,9,10-tetrahydro-imidazo-[1,2-h][1,7]naphthyridine

400 mg of the title compound of m.p. 235 - 236°C are obtained after crystallisation from isopropyl acetate by chromatographic purification on silica gel of the solid residue of Example 5.

Commercial utility

The compounds of the formula I and their salts have valuable pharmacological properties which make them commercially utilizable. In particular, they exhibit marked inhibition of gastric acid secretion and an excellent gastric and intestinal protective action in warm-blooded animals, in particular humans. In this connection, the compounds according to the invention are distinguished by a high selectivity of action, an advantageous duration of action, a particularly good enteral activity, the absence of significant side effects and a large therapeutic range.

"Gastric and intestinal protection" in this connection is understood as meaning the prevention and treatment of gastrointestinal diseases, in particular of gastrointestinal inflammatory diseases and lesions (such as, for example, gastric ulcer, duodenal ulcer, gastritis, hyperacidic or medicament-related functional dyspepsia), which can be caused, for example, by microorganisms (e.g. *Helicobacter pylori*), bacterial toxins, medicaments (e.g. certain antiinflammatories and antirheumatics), chemicals (e.g. ethanol), gastric acid or stress situations.

In their excellent properties, the compounds according to the invention surprisingly prove to be clearly superior to the compounds known from the prior art in various models in which the antiulcerogenic and the antisecretory properties are determined. On account of these properties, the compounds of the formula I and their pharmacologically acceptable salts are outstandingly suitable for use in human and veterinary medicine, where they are used, in particular, for the treatment and/or prophylaxis of disorders of the stomach and/or intestine.

A further subject of the invention are therefore the compounds according to the invention for use in the treatment and/or prophylaxis of the abovementioned diseases.

The invention likewise includes the use of the compounds according to the invention for the production of medicaments which are employed for the treatment and/or prophylaxis of the abovementioned diseases.

The invention furthermore includes the use of the compounds according to the invention for the treatment and/or prophylaxis of the abovementioned diseases.

A further subject of the invention are medicaments which comprise one or more compounds of the formula I and/or their pharmacologically acceptable salts.

The medicaments are prepared by processes which are known per se and familiar to the person skilled in the art. As medicaments, the pharmacologically active compounds according to the invention (= active compounds) are either employed as such, or preferably in combination with suitable pharmaceuti-

cal auxiliaries or excipients in the form of tablets, coated tablets, capsules, suppositories, patches (e.g. as TTS), emulsions, suspensions or solutions, the active compound content advantageously being between 0.1 and 95% and it being possible to obtain a pharmaceutical administration form exactly adapted to the active compound and/or to the desired onset of action (e.g. a sustained-release form or an enteric form) by means of the appropriate selection of the auxiliaries and excipients.

The auxiliaries and excipients which are suitable for the desired pharmaceutical formulations are known to the person skilled in the art on the basis of his/her expert knowledge. In addition to solvents, gel-forming agents, suppository bases, tablet auxiliaries and other active compound excipients, it is possible to use, for example, antioxidants, dispersants, emulsifiers, antifoams, flavor corrigents, preservatives, solubilizers, colorants or, in particular, permeation promoters and complexing agents (e.g. cyclodextrins).

The active compounds can be administered orally, parenterally or percutaneously.

In general, it has proven advantageous in human medicine to administer the active compound(s) in the case of oral administration in a daily dose of approximately 0.01 to approximately 20, preferably 0.05 to 5, in particular 0.1 to 1.5, mg/kg of body weight, if appropriate in the form of several, preferably 1 to 4, individual doses to achieve the desired result. In the case of a parenteral treatment, similar or (in particular in the case of the intravenous administration of the active compounds), as a rule, lower doses can be used. The establishment of the optimal dose and manner of administration of the active compounds necessary in each case can easily be carried out by any person skilled in the art on the basis of his/her expert knowledge.

If the compounds and/or their salts according to the invention are to be used for the treatment of the abovementioned diseases, the pharmaceutical preparations can also contain one or more pharmacologically active constituents of other groups of medicaments, for example: tranquilizers, (for example from the group of the benzodiazepines, for example diazepam), spasmolytics (for example, biefamiverine or camylofin), anticholinergics, (for example, oxyphencyclimine or phencarbamide), local anesthetics, (for example, tetracaine or procaine), and, if appropriate, also enzymes, vitamins or amino acids.

To be emphasized in this connection is in particular the combination of the compounds according to the invention with pharmaceuticals which inhibit acid secretion, such as, for example, H₂ blockers (e.g. cimetidine, ranitidine), H⁺/K⁺ ATPase inhibitors (e.g. omeprazole, pantoprazole), or further with so-called peripheral anticholinergics (e.g. pirenzepine, telenzepine) and with gastrin antagonists with the aim of increasing the principal action in an additive or super-additive sense and/or of eliminating or of decreasing the side effects, or further the combination with antibacterially active substances (such as, for example, cephalosporins, tetracyclines, penicillins, macrolides, nitroimidazoles or alternatively bis-

muth salts) for the control of *Helicobacter pylori*. Suitable antibacterial co-components which may be mentioned are, for example, mezlocillin, ampicillin, amoxicillin, cefalothin, cefoxitin, cefotaxim, imipenem, gentamycin, amikacin, erythromycin, ciprofloxacin, metronidazole, clarithromycin, azithromycin and combinations thereof (for example clarithromycin + metronidazole).

Pharmacology

The excellent gastric protective action and the gastric acid secretion-inhibiting action of the compounds according to the invention can be demonstrated in investigations on animal experimental models. The compounds according to the invention investigated in the model mentioned below have been provided with numbers which correspond to the numbers of these compounds in the examples.

Testing of the secretion-inhibiting action on the perfused rat stomach

In Table A which follows, the influence of the compounds according to the invention on the pentagastrin-stimulated acid secretion of the perfused rat stomach after intravenous administration in vivo is shown.

Table A

No.	Dose ($\mu\text{mol/kg}$) i.v.	Inhibition of acid secretion (%)
1	3	100
5	3	100

Methodology

The abdomen of anesthetized rats (CD rat, female, 200-250 g; 1.5 g/kg i.m. urethane) was opened after tracheotomy by a median upper abdominal incision and a PVC catheter was fixed transorally in the esophagus and another via the pylorus such that the ends of the tube just projected into the gastric lumen. The catheter leading from the pylorus led outwards into the right abdominal wall through a side opening.

After thorough rinsing (about 50-100 ml), warm (37°C) physiological NaCl solution was continuously passed through the stomach (0.5 ml/min, pH 6.8-6.9; Braun-Unita 1). The pH (pH meter 632, glass electrode EA 147; $\phi = 5$ mm, Metrohm) and, by titration with a freshly prepared 0.01 N NaOH solution to pH 7 (Dosimat 665 Metrohm), the secreted HCl were determined in the effluent in each case collected at an interval of 15 minutes.


The gastric secretion was stimulated by continuous infusion of 1 $\mu\text{g/kg}$ (= 1.65 ml/h) of i.v. pentagastrin (left femoral vein) about 30 min after the end of the operation (i.e. after determination of 2 preliminary

fractions). The substances to be tested were administered intravenously in a 1 ml/kg liquid volume 60 min after the start of the continuous pentagastrin infusion.

The body temperature of the animals was kept at a constant 37.8-38°C by infrared irradiation and heat pads (automatic, stepless control by means of a rectal temperature sensor).

Patent Claims

1. A compound of the formula I



in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl or hydroxy-1-4C-alkyl,

R3 is hydrogen or halogen,

one of the substituents R4a and R4b is hydrogen and the other is hydrogen, hydroxyl, 1-4C-alkoxy, 1-4C-alkoxy-1-4C-alkoxy, 1-4C-alkylcarbonyloxy or the radical R4', or R4a and R4b together are O (oxygen),

where,

R4' is -O-(CH₂)_m-S(O)_n-R₉,
 -S(O)_n-(CH₂)_m-OH,
 -S(O)_n-(CH₂)_m-O-R₉,
 -S(O)_n-(CH₂)_m-S(O)_p-R₉,
 -O-alk1-S(O)_n-R₉,
 -S(O)_n-R₉,
 -S(O)_n-alk1-OH,
 -S(O)_n-alk1-O-R₉ or
 -S(O)_n-alk1-S(O)_p-R₉,

in which

R₉ is 1-7C-alkyl, halo-1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl or di-1-4C-alkylamino-1-4C-alkyl, Ar or Ar-1-4C-alkyl, where Ar is phenyl or substituted phenyl having one, two or three identical or different substituents selected from the group consisting of 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkylcarbonyl,

1-4C-alkoxycarbonyl, halogen, trifluoromethyl, difluoromethoxy, trifluoromethoxy, amino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino and nitro,

alk1 is 1-4C-alkyl-, hydroxyl-, oxo-, carboxyl-, halogen-, amino-, 1-4C-alkoxy-carbonylamino- or phenyl-substituted 2-7C-alkylene or 3-4C-alkenylene,

m is an integer from 2 to 7,

n is the number 0, 1 or 2 and

p is the number 0, 1 or 2,

one of the substituents R5a and R5b is hydrogen and the other is hydrogen, hydroxyl, 1-4C-alkoxy, 1-4C-alkoxy-1-4C-alkoxy, 1-4C-alkylcarbonyloxy or the radical R5', or R5a and R5b together are O (oxygen)

where

R5' is -O-(CH₂)_q-S(O)_r-R10,
 -S(O)_r-(CH₂)_q-OH,
 -S(O)_r-(CH₂)_q-O-R10,
 -S(O)_r-(CH₂)_q-S(O)_t-R10,
 -O-alk2-S(O)_r-R10,
 -S(O)_r-R10,
 -S(O)_r-alk2-OH,
 -S(O)_r-alk2-O-R10 or
 -S(O)_r-alk2-S(O)_t-R10,

in which

R10 is 1-7C-alkyl, halo-1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl or di-1-4C-alkylamino-1-4C-alkyl, Ar or Ar-1-4C-alkyl, where Ar is phenyl or substituted phenyl having one, two or three identical or different substituents selected from the group consisting of 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkylcarbonyl, 1-4C-alkoxy-carbonyl, halogen, trifluoromethyl, difluoromethoxy, trifluoromethoxy, amino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino and nitro,

alk2 is 1-4C-alkyl-, hydroxyl-, oxo-, carboxyl-, halogen-, amino-, 1-4C-alkoxycarbonylamino- or phenyl-substituted 2-7C-alkylene or 3-4C-alkenylene,

q is an integer from 2 to 7,

r is the number 0, 1 or 2 and

t is the number 0, 1 or 2,

R6 is hydrogen, halogen, 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or trifluoromethyl,


R7 is hydrogen, halogen, 1-4C-alkyl or 1-4C-alkoxy and

R8 is hydrogen or 1-4C-alkyl,

and its salts,

where one of the substituents R4a and R4b has to be R4' and/or one of the substituents R5a and R5b has to be R5'.

2. A compound as claimed in claim 1, of the formula I*



in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl or hydroxy-1-4C-alkyl,

R3 is hydrogen,

one of the substituents R4a and R4b is hydrogen and the other is hydrogen, hydroxyl, 1-4C-alkoxy, 1-4C-alkoxy-1-4C-alkoxy or the radical R4',

where

R4' is -O-(CH₂)_m-S(O)_n-R9,
 -S(O)_n-(CH₂)_m-OH,
 -S(O)_n-(CH₂)_m-O-R9,
 -S(O)_n-(CH₂)_m-S(O)_p-R9 or
 -S(O)_n-R9,

in which

R9 is 1-7C-alkyl, halo-1-4C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

m is an integer from 2 to 7,


n is the number 0, 1 or 2 and

p is the number 0, 1 or 2,

one of the substituents R5a and R5b is hydrogen and the other is hydrogen, hydroxyl, 1-4C-alkoxy, 1-4C-alkoxy-1-4C-alkoxy or the radical R5',

where

R5' is -O-(CH₂)_q-S(O)_r-R10,
 -S(O)_r-(CH₂)_q-OH,
 -S(O)_r-(CH₂)_q-O-R10,



in which

R10 is 1-7C-alkyl, halo-1-4C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

q is an integer from 2 to 7,

r is the number 0, 1 or 2 and

t is the number 0, 1 or 2,

R6 is hydrogen, halogen or trifluoromethyl,

R7 is hydrogen or halogen and

R8 is hydrogen,

and its salts,

where one of the substituents R4a and R4b has to be R4' and/or one of the substituents R5a and R5b has to be R5'.

3. A compound I* as claimed in claim 2 in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl,

R3 is hydrogen,

one of the substituents R4a and R4b is hydrogen and the other is R4'

where

R4' is $-\text{O}-(\text{CH}_2)_m-\text{S(O)}_n\text{-R9}$,

$-\text{S(O)}_n\text{-R9}$,

$-\text{S(O)}_n\text{-(CH}_2\text{)}_m\text{-OH}$ or

$-\text{S(O)}_n\text{-(CH}_2\text{)}_m\text{-O-R9}$,

in which

R9 is 1-4C-alkyl,

m is the number 2 or 3 and

n is the number 0, 1 or 2,

one of the substituents R5a and R5b is hydrogen and the other is hydroxyl,

R6 is hydrogen,

R7 is hydrogen and

R8 is hydrogen,

and its salts.

4. A compound I* as claimed in claim 2 in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl,

R3 is hydrogen,

one of the substituents R4a and R4b is hydrogen and the other is R4',

where

R4' is -O-(CH₂)_m-S(O)_n-R9,
 -S(O)_n-R9,
 -S(O)_n-(CH₂)_m-OH or
 -S(O)_n-(CH₂)_m-O-R9,

in which

R9 is 1-4C-alkyl,

m is the number 2 or 3 and

n is the number 0, 1 or 2,

one of the substituents R5a and R5b is hydrogen and the other is 1-4C-alkoxy or 1-4C-alkoxy-1-4C-alkoxy,

R6 is hydrogen,

R7 is hydrogen and

R8 is hydrogen,

and its salts.

5. A compound I* as claimed in claim 2 in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl,

R3 is hydrogen,

one of the substituents R4a and R4b is hydrogen and the other is R4',

where

R4' is -O-(CH₂)_m-S(O)_n-R9,
 -S(O)_n-R9,
 -S(O)_n-(CH₂)_m-OH or
 -S(O)_n-(CH₂)_m-O-R9,

in which

R9 is 1-4C-alkyl,

m is the number 2 or 3 and

n is the number 0, 1 or 2,

one of the substituents R5a and R5b is hydrogen and the other is R5',

where

R5' is -O-(CH₂)_q-S(O)_r-R10,
 -S(O)_r-R10,
 -S(O)_r-(CH₂)_q-OH or
 -S(O)_r-(CH₂)_q-O-R10,

in which

R10 is 1-4C-alkyl,

q is the number 2 or 3 and

r is the number 0, 1 or 2,

R6 is hydrogen,

R7 is hydrogen and

R8 is hydrogen,

and its salts.

6. A compound I* as claimed in claim 2 in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl,

R3 is hydrogen,

one of the substituents R4a and R4b is hydrogen and the other is 1-4C-alkoxy or 1-4C-alkoxy-1-4C-alkoxy,

one of the substituents R5a and R5b is hydrogen and the other is R5',

where

R5' is $-O-(CH_2)_q-S(O)_r-R10$,

$-S(O)_r-R10$,

$-S(O)_r-(CH_2)_q-OH$ or

$-S(O)_r-(CH_2)_q-O-R10$,

in which

R10 is 1-4C-alkyl,

q is the number 2 or 3 and

r is the number 0, 1 or 2,

R6 is hydrogen,

R7 is hydrogen and

R8 is hydrogen,

and its salts.

7. A compound I* as claimed in any of claims 2, 3, 4, 5 or 6 in which R5b is hydrogen.

8. A compound I* as claimed in any of claims 2, 3, 4, 5 or 6 in which R5b is hydrogen and R4a is hydrogen.

9. A medicament comprising a compound as claimed in claim 1 and/or a pharmacologically acceptable salt thereof together with customary pharmaceutical auxiliaries and/or excipients.

10. The use of compounds as claimed in claim 1 and their pharmacologically acceptable salts for preparing medicaments for the prevention and treatment of gastrointestinal disorders.

International Application No.

PCT/EP 99/08227

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D471/14 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 42707 A (BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH) 1 October 1998 (1998-10-01) cited in the application claims 1,3,9,10	1,2,9,10
Y	US 4 468 400 A (E. H. GOLD ET AL.) 28 August 1984 (1984-08-28) cited in the application claims 1,12	1,2,9,10
A	EP 0 235 575 A (G. D. SEARLE & CO.) 9 September 1987 (1987-09-09) cited in the application	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

*** Special categories of cited documents:****"A"** document defining the general state of the art which is not considered to be of particular relevance**"E"** earlier document but published on or after the international filing date**"L"** document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)**"O"** document referring to an oral disclosure, use, exhibition or other means**"P"** document published prior to the international filing date but later than the priority date claimed**"T"** later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention**"X"** document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone**"Y"** document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art**"&"** document member of the same patent family

Date of the actual completion of the international search

17 February 2000

Date of mailing of the international search report

24/02/2000

Name and mailing address of the ISA

European Patent Office, P.B. 6818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hass, C

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/08227

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9842707	A	01-10-1998	AU 7520898 A EP 0971922 A HR 980147 A NO 994584 A	20-10-1998 19-01-2000 28-02-1999 23-11-1999
US 4468400	A	28-08-1984	NONE	
EP 235575	A	09-09-1987	US 4755524 A US 4711903 A AT 56702 T AU 593606 B AU 6810387 A CA 1302424 A DK 50487 A FI 870417 A,B, GR 3000864 T JP 2089898 C JP 8005852 B JP 62228054 A KR 9102948 B NO 870384 A,B, NZ 219104 A PH 24624 A PT 84229 A,B ZA 8700657 A	05-07-1988 08-12-1987 15-10-1990 15-02-1990 06-08-1987 02-06-1992 01-08-1987 01-08-1987 15-11-1991 02-09-1996 24-01-1996 06-10-1987 11-05-1991 03-08-1987 27-09-1989 17-08-1990 01-02-1988 30-03-1988